

# Consume the electrolyte solution 20 minutes before physical exercise reduce systolic blood pressure and pulse rate

*by I Wayan Teresna*

---

**Submission date:** 09-May-2023 12:37PM (UTC+0700)

**Submission ID:** 2088317798

**File name:** GSCBPS-2022-0308.pdf (798.37K)

**Word count:** 3699

**Character count:** 18578



(RESEARCH ARTICLE)



## Consume the electrolyte solution 20 minutes before physical exercise reduce systolic blood pressure and pulse rate

I Nengah Sandi <sup>1,\*</sup>, Agustinus Dei <sup>2</sup>, Daniel Womsiwor <sup>3</sup> and I Wayan Teresna <sup>4</sup>

<sup>1</sup> Department of Physics, Faculty of Mathematics and Natural Sciences, Udayana University, Bali, Indonesia.

<sup>2</sup> Department of Physical Education, Health and Recreation, Faculty of Teacher Training and Education, PGRI Mahadewa University, Indonesia.

<sup>3</sup> Department of Training, Faculty of Sports Science, Cenderawasih University Papua, Indonesia.

<sup>4</sup> Department of Electrical Engineering, Bali State Polytechnic, Indonesia.

GSC Biological and Pharmaceutical Sciences, 2022, 20(01), 340-345

Publication history: Received on 26 May 2022; revised on 31 July 2022; accepted on 02 August 2022

Article DOI: <https://doi.org/10.30574/gscbps.2022.20.1.0308>

### Abstract

Physical exercise can increase systolic blood pressure (SBP), diastolic blood pressure (DBP), pulse rate (FR), and body temperature (BT) exercise. The increase in the body's vital signs can be overcome by consuming electrolyte solutions before exercise. The purpose of this study was to determine the effect of consuming 500 mL of electrolyte solution orally 20 minutes before exercise. The intensity of exercise is 70% of the maximum pulse rate with a duration of 30 minutes, the effect of which is to decrease SBP, DBP, PR, and exercise BT. This experimental study used a pre and post test control group design, in which 18 subjects were divided into two groups and given different treatments, namely Klp-1 was given mineral water and Klp-2 was given electrolyte solution with the same volume. Differences in treatment results were analyzed using the T-Independent test at the significance of  $< 0.05$ . The difference in mean SBP, DBP, PR, and BT between groups after exercise was  $p = 0.005$ ,  $p = 0.320$ ,  $p = 0.000$ , and  $p = 0.100$ . So giving 500 mL of electrolyte solution 20 minutes before physical exercise can reduce SBP and PR ( $p < 0.05$ ) but can't reduce DBP and BT ( $p > 0.05$ ). Therefore, it is recommended to consume 500 mL of electrolyte solution 20 minutes before physical exercise.

**Keywords:** Electrolyte solution; Systolic blood pressure; Diastolic blood pressure; Pulse rate

### 1. Introduction

Sport is a planned, structured physical activity that is performed repetitively. The goal of the sport is to improve or maintain physical fitness. Therefore, exercise is an important thing that must be done continuously for a long time.

Exercise has many benefits for the human body, such as making the body fit and stamina is always maintained. In addition, exercise can also burn body fat when done for a long duration [1]. Sports or physical activity has the potential to increase PR if the activity load is increased. This happens because the increased physical activity of the body, the more blood flow to supply nutrients and oxygen to active tissues, so that the heart contracts faster and stronger. This increase also causes an increase in PR [2].

It is also stated that dynamic physical exercise involving large muscles will have an effect on increasing cardiovascular function, namely SBP, DBP, and PR, as well as increasing cardiac output [3]. In addition, every physical exercise the body always produces heat which depends on the intensity and duration of the exercise. The higher the activity, the higher the body weight produced and the longer the duration of the exercise, the higher the body temperature [4].

\*Corresponding author: I Nengah Sandi

Department of Physics, Faculty of Mathematics and Natural Sciences, Udayana University, Bali, Indonesia.

Copyright © 2022 Author(s) retain the copyright of this article. This article is published under the terms of the Creative Commons Attribution License 4.0.

Without realizing it, after doing activities for a long time, the body loses a lot of fluids. This fluid loss makes you feel thirsty and if not replaced immediately, will lead to dehydration. Dehydration is a decrease in body fluids due to sweating without being replaced with fluids through drinking in moderation [5]. Excretion of body fluids by means of sweating aims to release body heat due to increased activity. The release of body heat aims to stabilize the body's core temperature. The ultimate goal is to minimize the risk of heat injury [6].

The human body is made up of mostly fluids. Water and electrolytes contained in body fluids are needed for the continued effectiveness of nerves and muscles. Heavy and prolonged physical exercise results in a buildup of lactic acid and an increase in body fluid expenditure through sweat. During physical activity, body fluids are needed to maintain balance and the body's metabolic processes. If fluid intake is lower than expenditure, the body will experience a fluid balance disorder called dehydration. Electrolytes are chemical compounds in solution that dissociate into positively or negatively charged particles. Most of the metabolism requires and is influenced by electrolytes. The major contents of electrolytes in body fluids that are needed by the body are sodium (Na<sup>+</sup>), potassium (K<sup>+</sup>), chloride (Cl<sup>-</sup>), and bicarbonate (HCO<sub>3</sub><sup>-</sup>) [7].

## 1 2. Material and methods

This research is an experimental study using a pre and post-test control group design. The research was carried out at Ngurah Rai Bali Sports Center in the morning at 07.00-09.00. The number of samples in this study were 18 students who had met the inclusion criteria, namely aged 16-18 years, were willing to participate as research subjects and filled out informed consent. The exclusion criteria were if the participant was sick based on a doctor's examination. Participants will be excluded from the study if they withdraw without a reason. Sample selection and sample allocation were carried out using a simple random sampling technique, so that each group consisted of 9 people.

Physical exercise was carried out for 30 minutes on the control group and the treatment group on different days. Before starting the exercise, blood pressure, pulse frequency, and body temperature were measured before treatment, and after that warm-up was done. Body weight and height were also measured to determine body mass index before treatment. The control group was given 500 mL of mineral water and the treatment group was given the same amount of electrolyte fluid, 20 minutes before the start of the exercise. Immediately after the exercise is complete, blood pressure, pulse frequency, and body temperature will be measured again. After that, the body mass index was measured again after treatment.

34  
The data obtained were analyzed using the Saphiro Wilk test to test for normality, the Levene Test to test the homogeneity of the data, and the Independent T-Test to determine the difference in treatment results between groups.

## 3. Results

The research data concerns the characteristics of research subjects, data on the research environment, and research results.

2  
Table 1 Physical Characteristics of Research Subjects

Variable	Control Group			Treatment Group		
	Minimum	Maximum	Mean	Minimum	Maximum	Mean
Age (years)	17.0	18.0	17.4	17.0	17.0	17.0
Weight (kg)	54.1	70.3	61.6	51.2	74.0	64.9
Height (cm)	166.0	178.0	170.2	159.0	176.0	169.8
Body Mass Index (kg/m <sup>2</sup> )	18.6	23.5	21.2	18.6	24.8	22.5

The characteristics data of the research results taken before and after treatment are presented as in Table 3. The data measured were systolic blood pressure, diastolic blood pressure, pulse frequency and body temperature.

2  
Before determining the statistical test used, the normality and homogeneity of the data were first tested. The data tested were TDS, TDD, FDN, and BT both before and after treatment (Table 4).

**Table 2** Characteristics of the Research Environment

Variable	Minimum	Maximum	Mean
Ambient Temperature before Treatment	28.2	28.8	28.5
Ambient Temperature after Treatment	31.0	31.4	31.2
Relative Humidity before Treatment	56.0	59.0	57.5
Relative Humidity after Treatment	45.0	47.0	46.0

**Table 3** Characteristics of Research Results

Variable	Control Group			Treatment Group		
	Minimum	Maximum	Mean	Minimum	Maximum	Mean
<b>Before Treatment</b>						
SBP (mmHg)	112.0	138.0	127.1	106.0	128.0	115.9
DBP (mmHg)	65.0	87.0	75.7	63.0	82.0	74.6
PR (beats/minute)	73.0	97.0	82.7	63.0	76.0	70.4
BT (°C)	36.5	37.5	36.9	36.3	37.1	36.6
<b>After Treatment</b>						
SBP (mmHg)	134.0	158.0	145.2	130.0	142.0	135.6
DBP (mmHg)	88.0	99.0	93.4	85.0	96.0	89.4
PR (beats/minute)	110.0	134.0	121.8	96.0	105.0	100.4
BT (°C)	37.1	37.8	37.4	36.5	37.6	36.9

Information: SBP = systolic blood pressure, DBP = diastolic blood pressure, PR = pulse rate, BT = body temperature

**Table 4** Normality and Data Homogeneity Test Results

Variable	p-Value Normality (Saphiro Wilk)		p-Value Homogeneity (Levene Test)
	Control Group	Treatment Group	
SBP before Treatment	0.510	0.483	0.621
SBP after Treatment	0.888	0.992	0.123
DBP before Treatment	0.612	0.215	0.819
DBP after Treatment	0.731	0.172	0.378
PR before Treatment	0.405	0.685	0.092
PR after Treatment	0.908	0.555	0.052
BT before Treatment	0.828	0.231	0.649
BT after Treatment	0.477	0.634	0.178

Information: SBP = systolic blood pressure, DBP = diastolic blood pressure, PR = pulse rate, BT = body temperature

SBP, DBP, PR, and BT before and after treatment were normally distributed ( $P > 0.05$ ). The data variance between groups both before and after treatment was homogeneous ( $P > 0.05$ ). Because the variable data being compared were normally

distributed and homogeneous, the mean difference in the data after treatment was analyzed using the Independent T-Test.

Because the data before treatment between groups was homogeneous, it can be stated that the initial state of the data came from the same source, so that the difference in SBP, DBP, PR, and BT after treatment was due to differences in the treatment given. To determine the effect of treatment on SBP, DBP, PR, and BT after treatment, it is presented in Table 5.

**Table 5** Results of the Mean Difference between Groups After Treatment

Variable	Mean of Control Group	Mean of Treatment Group	Mean Difference	p-Value
SBP	145.2	135.6	9.6	0.005
DBP	93.4	89.4	4.0	0.320
PR	121.8	100.4	21.4	0.000
BT	37.4	36.9	0.5	0.100

Information: SBP = systolic blood pressure, DBP = diastolic blood pressure, PR = pulse rate, BT = body temperature

The difference in mean SBP and PR between groups after treatment was significantly different with a value of  $p < 0.05$ , while the mean difference between DBP and BT after treatment was not significantly different with a value of  $p > 0.05$ .

#### 4. Discussion

Table 1 shows that the mean age in the two groups is almost the same age, which is between 17-18 years. This is in accordance with the age of high school students, which ranges from 16-18 years. The effect of age on resting pulse rate can be seen from the resting pulse rate in adults as much as 60-90 beats per minute [8]. The results of previous research on 33 students of SMKN-5 Denpasar, it was found that there was a significant relationship between age and resting pulse rate with  $p = 0.007$  ( $p < 0.05$ ) [9].

Body weight and height are used to determine body mass index. Based on the characteristics of body mass index, it shows that the minimum value in the control group is  $18.6 \text{ kg/m}^2$  and the treatment group is  $18.10 \text{ kg/m}^2$ . The maximum value in the control group was  $23.5 \text{ kg/m}^2$  and the treatment group was  $24.8 \text{ kg/m}^2$ . The mean in the control group was  $21.2 \text{ kg/m}^2$  and the treatment group was  $22.5 \text{ kg/m}^2$ . Judging from the subject's body mass index, this value is within the normal limits of WHO standards [10]. So based on the body mass index, the nutritional status of the subject was within normal limits, thus the subject was not overweight or obese.

Table 2 shows that the environmental characteristics of the control group and the treatment group consisted of ambient temperature and relative humidity of the air. The minimum value of the ambient temperature of the two groups before treatment was  $28.2^\circ\text{C}$  and the maximum was  $28.8^\circ\text{C}$ , while the mean value was  $28.5^\circ\text{C}$ . The minimum value of the environmental temperature of the two groups after treatment was  $31.0^\circ\text{C}$  and the maximum value was  $31.4^\circ\text{C}$ , while the mean value was  $31.2^\circ\text{C}$ . The relative humidity of the air before treatment showed that the minimum value for both groups was 56.0% and the maximum value was 59.0%, while the average was 57.5%. The minimum value for both groups after treatment was 45.0% and the maximum value was 47.0%, while the mean value was 46.0%. The results of previous studies showed that 40% relative humidity is better than 50% and 60% humidity because it can inhibit the increase in pulse rate, body temperature and exercise blood lactic acid [11]. This value indicates that the relative humidity is within normal limits. So that the subject can perform physical activity comfortably. The WBGT Index (Wet Bulb Globe Temperature Index) is used to assess the level of environmental comfort that combines the impact of solar radiation, ambient temperature, humidity and wind speed. In various sports the WBGT index is recommended to reach  $25^\circ\text{C}$ , and exercise is considered unsafe if the WBGT index reaches more than  $28^\circ\text{C}$  for those who are not trained or have not acclimatized. This value is in the normal to high limit [12].

According to Table 3, the mean SBP before treatment in the control group was 127.1 mmHg and the treatment group was 115.9 mmHg. The mean DBP of the control group was 75.7 mmHg and the treatment group was 74.6 mmHg. The normal state of resting SBP is in the range of 110.0-120.0 mmHg while DBP is between 70.0-80.0 mmHg. So that the SBP of subjects in the control group slightly increased and the treatment group was within normal limits, while the SBP of subjects in the control group and the treatment group was within normal limits.



The mean PR of the control group was 82.7 beats/minute and the treatment group was 70.4 beats/minute. So that the resting pulse frequency of the research subjects was within normal limits. The effect of age on the resting pulse rate can be seen from the resting pulse rate in adults as much as 60-80 beats/minute. This statement is in accordance with the results of research on 33 students of UN-5 Denpasar, it was found that there was a significant relationship between age and resting pulse frequency with  $p = 0.007$  ( $p < 0.05$ ) [9].

The mean ST in the control group was 36.9°C and the treatment group was 36.6°C. Normal body temperature is in the range of 36.3°C-37.1°C. So that the ST research subjects are in normal limits.

The mean SBP after treatment in the control group was 145.2 mmHg and the treatment group was 135.6 mmHg. Systolic blood pressure at rest 120 mmHg will increase to 140-250 mmHg during maximum intensity exercise. This is possible because most of the skeletal muscles contract which forces the blood vessels. The increased pressure only lasts a few minutes, and the pressure will slowly return to normal. This increase was also followed by DBP, but the increase was not too high [4]. So that the increase in SBP after treatment of the subject was within normal limits.

The mean PR after treatment in the control group was 121.8 beats/minute and the treatment group was 100.4 beats/minute. Exercise PR will increase up to 185 beats / min, where the increase reaches 270% which will increase cardiac output [13]. The increase in exercise PR in research subjects is within normal limits. It is also stated that physical activity causes acute changes in respiratory rate and PR [14].

The mean ST after treatment in the control group was 37.4°C and the treatment group was 36.9°C. During physical activity, ST will increase up to 40°C [13]. So that the increase in ST in this research subject is still within normal limits.

Table 5 shows that there was a significant difference in SBP and PR exercise between the control group compared to the treatment group, where SBP and PR exercise in the control group were higher than the treatment group. This difference was caused by differences in the treatment given, namely before the treatment the control group was only given drinking water while the control group was given drinking electrolyte fluids.

Drinking water is recommended for someone who is active for a relatively short period of time (<75 minutes), but electrolyte drinks containing carbohydrates are highly recommended and are the average choice for athletes [15]. The research subjects were given 500 ml of electrolyte fluid in the form of Pocari Sweat with a total fat content of 0 g, cholesterol 0 g, sodium 125 mg, total carbohydrates 15 g, and protein 0 g. Administration of electrolyte-carbohydrate fluids can reduce the level of dehydration and reduce the effect of fluid loss on cardiovascular function [16].

The purpose of providing electrolyte-carbohydrate fluids is to stimulate the rate of fluid absorption, supply of carbohydrates and other nutrients as a substrate for physical activity, increase rehydration, decrease physiological stress and recovery after exercise. Maintaining fluid levels of carbohydrates is very important, 1% of fluid is lost from sweat which will eventually result in decreased exercise performance. According to the research results of Hornsby [17], electrolyte-carbohydrate drinks can reduce pulse frequency, systolic blood pressure and diastolic blood pressure.

Water is an essential ingredient in every life, which is the main key to survival, namely by preventing dehydration. Without water humans can only survive for a few days. When dehydration occurs, water is a temporary way to replace the lost fluids. Subsequent replacement is with electrolyte fluids for moderate to severe dehydration [18].

So because the drinks consumed by the subjects in the treatment group were electrolyte fluids with a total fat content of 0 g, cholesterol 0 g, sodium 125 mg, total carbohydrates 15 g, protein 0 g, the electrolyte fluid used in this study could reduce SBP and PR. according to the results of previous research.

## 5. Conclusion

Consuming 500 mL of electrolyte solution 20 minutes before exercise with an intensity of 70% maximum pulse rate can reduce exercise SBP by 6.6% and exercise PR by 17.6% ( $p < 0.05$ ). Besides that, it can also reduce DBP by 4.3% and exercise BT by 1.3%, but the decrease is not significantly different ( $p > 0.05$ ). Consuming 500 mL of electrolyte solution before physical exercise can't significantly reduce diastolic blood pressure and exercise body temperature, therefore it is necessary to re-examine why it occurs.

12

## Compliance with ethical standards

### Acknowledgments

With the success of this research, we would like to express our deepest gratitude to the Head of SMA-5 Denpasar, sports teacher, and the manager of the Gusti Ngurah Rai Bali Sports Center for lending a place. We would also like to thank our friends who have helped the smooth running of this research and all students who have taken their time as research subjects.

12

### Disclosure of conflict of interest

For the sake of the smooth running of the publication, we as the authors agree to provide flexibility to the correspondent writers as the main authors to arrange the writing format and regulate the publishing costs in accordance with the publisher's provisions.

### Statement of informed consent.

No informed consent.

## References

- [1] Fatmah. Fitness and Sports Nutrition. Bandung: Lubuk Agung. 2011.
- [2] Ganong WF. 2012. Review of Medical Physiology, 22 Ed. California: McGraw-Hill Companies, Inc. 2012.
- [3] Kadir A. Cardiovascular Adaptation to Physical Exercise. [cited 2016 March, 27]. Available from: [RL://fk.uwks.ac.id/jurnal/daftar\\_edisi](http://fk.uwks.ac.id/jurnal/daftar_edisi). 2010.
- [4] Sandi N. Effect of Temperature and Relative Humidity of Air on Physical Appearance in Sports. National Seminar on Integration of Biodiversity and Culture in Sustainable Development. Denpasar 27 Nopember. 2014.
- [5] Sandi N, Parwata MY. Sports Environment. Yogyakarta: Wahana Rwsolusi. 2018.
- [6] Sandi N, Ariyasa G, Teresna W, Ashadi K. Effect of Relative Humidity on Changes in Body Temperature Exercise. Sport and Fitness Journal, 2017, 5(1): 103-109.
- [7] Hamidin AS. The goodness of white water. Editor: Retino. 1 Mold. Yogyakarta: Media Pressindo. 2010.
- [8] Burke ER. Complete Home Fitness Handbook. Washington: Human Kinetics. 2001.
- [9] Sandi N. The Relationship Between Height, Weight, Body Mass Index, and Age Against Resting Heart Rate, Students of SMKN 5 Denpasar. Sport and Fitness Journal. 2013, 1(1): 38-44.
- [10] Soetjningsih. Child Growth. Jakarta: EGC. 2013.
- [11] Sandi N, Adiputra N, Pangkahila A, Adiatmika PG. Relative Humidity of 40% Inhibiting the Increase of Pulse Rate, Body Temperature, And Blood Lactic Acid During Exercise. Bali Medical Journal. 2016, 5(2): 30-34.
- [12] Sandi N. Influence of Physical Activity on Blood Pressure. Joint International Conference APCHI-ERGOFUTURE-PEI-IAIFI. Denpasar 22-25 October. 2014.
- [13] Hall JE. Guyton and Hall Textbook of Medical Physiology (13th Edition). Philadelphia: John F. Kennedy Blvd. 2016.
- [14] Sandi N. Effect of Physical Exercise on Pulse Rate. Sport and Fitness Journal. 2016, 4(2): 1-6.
- [15] Kalman D, Feldman S. Comparison of Coconut Water and A Carbohydrate-Electrolyte Sport Drink on Measures of Hydration and Physical Performance In-Exercise Trained Men. Memphis: The University of Memphis, Cardiorespiratory / Metabolic Laboratory, 2012.
- [16] Ostojic SM, Mazic S. Effect of A Carbohydrate – Electrolyte Drink on Specific Soccer Tests and Performance. Journal of Sports Science and Medicine. 2002, 1: 47-53.
- [17] Hornsby J. The Effect of Carbohydrate-Electrolyte Sports Drinks on Performance and Physiological Function During An 8km Cycle Time Trial. Plymouth: School of Biomedical & Biological Sciences. 2011.
- [18] Popkin BM, Ancu KED, Rosenberg IH. Water, Hydration and Health. NIH Public Access, 2011, 68(8): 439-458.

Consume the electrolyte solution 20 minutes before physical exercise reduce systolic blood pressure and pulse rate

ORIGINALITY REPORT

17%

SIMILARITY INDEX

13%

INTERNET SOURCES

10%

PUBLICATIONS

6%

STUDENT PAPERS

PRIMARY SOURCES

1	Submitted to Universitas Airlangga Student Paper	2%
2	erepo.unud.ac.id Internet Source	1%
3	mdpi-res.com Internet Source	1%
4	Submitted to Udayana University Student Paper	1%
5	Submitted to Universitas Negeri Jakarta Student Paper	1%
6	ebin.pub Internet Source	1%
7	I MADE MULIARTA, I NYOMAN ADIPUTRA, I MADE KRISNA DINATA, LUH MADE INDAH S.H.A, I KETUT TUNAS. "ACTIVE STRETCHING AND WORKING POSTURE CORRECTION TO IMPROVE PSYCHO-PHYSIOLOGICAL RESPONSE AMONG COMPUTER OPERATORS	1%



FOR HIGH SCHOOL STUDENTS", Journal of  
Human Ergology, 2020

Publication

8

[researchoutput.csu.edu.au](https://researchoutput.csu.edu.au)

Internet Source

1 %

9

Tapanee Hongratanaworakit. "Aroma-  
therapeutic Effects of Massage Blended  
Essential Oils on Humans", Natural Product  
Communications, 2011

Publication

1 %

10

Justyna Drzał-Grabiec, Sławomir Snela, Maciej  
Rachwał, Justyna Rykała, Justyna Podgórska.  
"Effects of carrying a backpack in a  
symmetrical manner on the shape of the  
feet", Ergonomics, 2013

Publication

1 %

11

Hakan Doneray, Ozlem Ziraatci Akbulut, Ayse  
Ozden, Abdulkadir Yildirim, Zerrin Orbak.  
"Plasma renin, aldosterone, and urinary  
prostaglandin E2 levels in children with  
hypocalcemia due to vitamin D deficiency  
rickets", Prostaglandins & Other Lipid  
Mediators, 2022

Publication

<1 %

12

[magnascientiapub.com](https://magnascientiapub.com)

Internet Source

<1 %

13

[www.ncbi.nlm.nih.gov](https://www.ncbi.nlm.nih.gov)

Internet Source

<1 %

14

Tsai, W.K.. "Pigtail catheters vs large-bore chest tubes for management of secondary spontaneous pneumothoraces in adults", American Journal of Emergency Medicine, 200611

Publication

<1 %

15

N S Harahap, S Suprayitno, N Simatupang, R M Sari. "Lactic acid level in soccer who consume red dragon fruit juice and regular exercise", Journal of Physics: Conference Series, 2021

Publication

<1 %

16

Submitted to University of Aberdeen

Student Paper

<1 %

17

[article.imrpress.com](http://article.imrpress.com)

Internet Source

<1 %

18

[bmcpulmed.biomedcentral.com](http://bmcpulmed.biomedcentral.com)

Internet Source

<1 %

19

[www.karger.com](http://www.karger.com)

Internet Source

<1 %

20

[core.ac.uk](http://core.ac.uk)

Internet Source

<1 %

21

[zenodo.org](http://zenodo.org)

Internet Source

<1 %

22	<a href="http://editorialexpress.com">editorialexpress.com</a> Internet Source	<1 %
23	<a href="http://issuu.com">issuu.com</a> Internet Source	<1 %
24	<a href="http://jssm.org">jssm.org</a> Internet Source	<1 %
25	<a href="http://jurnal.ugm.ac.id">jurnal.ugm.ac.id</a> Internet Source	<1 %
26	<a href="http://www.nepjol.info">www.nepjol.info</a> Internet Source	<1 %
27	<a href="http://gjournals.org">gjournals.org</a> Internet Source	<1 %
28	<a href="http://link.springer.com">link.springer.com</a> Internet Source	<1 %
29	<a href="http://mattioli1885journals.com">mattioli1885journals.com</a> Internet Source	<1 %
30	<a href="http://pearl.plymouth.ac.uk">pearl.plymouth.ac.uk</a> Internet Source	<1 %
31	<a href="http://www.researchsquare.com">www.researchsquare.com</a> Internet Source	<1 %
32	<a href="http://www.science.gov">www.science.gov</a> Internet Source	<1 %
33	"Abstracts", Public Health Nutrition, 2013 Publication	<1 %

34

Novita Sari Harahap, Nurhayati Simatupang, Suprayitno Suprayitno. "The Effect of Physical Activity and Red Dragon Fruit (*Hylocereus polyrhizus*) in Red Blood Cell and Hemoglobin in Trained People", Open Access Macedonian Journal of Medical Sciences, 2021

Publication

<1 %

---

Exclude quotes  On

Exclude matches  < 3 words

Exclude bibliography  On